

**Problem Set 4**

February 27, 2023

**due: March 13, 2023**

All numbered exercises are from the textbook *Real Analysis, Foundations and Functions of One Variable*, by Laczkovich and Sos.

1. Let  $(a_n)_{n=1}^\infty$  be a sequence defined recursively as follows:  $a_1 = \sqrt{2}$ , and  $a_{n+1} = \sqrt{2 + a_n}$ , for all  $n \geq 1$ . Prove that the sequence converges and find its limit.
2. Let  $(a_n)_{n=1}^\infty$  be a bounded sequence, and let  $S$  denote the set of all *subsequential limits* of  $(a_n)_{n=1}^\infty$ , that is, all real numbers  $s$  such that there exists a subsequence  $(a_{n_k})_{k=1}^\infty$  of  $(a_n)_{n=1}^\infty$  with  $\lim_{k \rightarrow \infty} a_{n_k} = s$ . One defines *limit superior* of  $(a_n)_{n=1}^\infty$ , denoted  $\limsup a_n$ , as  $\sup S$ , and *limit inferior* of  $(a_n)_{n=1}^\infty$ , denoted  $\liminf a_n$ , as  $\inf S$ . Prove the following:

(a)  $\limsup a_n = \lim_{N \rightarrow \infty} \sup\{a_n : n \geq N\}$

(b)  $\liminf a_n = \lim_{N \rightarrow \infty} \inf\{a_n : n \geq N\}$

(c) For every  $s_0 \in \mathbb{R}$ , if there exists a sequence  $(s_k)_{k=1}^\infty$  with values in  $S$ , such that  $\lim_{k \rightarrow \infty} s_k = s_0$ , then  $s_0 \in S$ .

3. **Construction of  $\mathbb{R}$ :** Let  $\mathbb{Q}$  denote the ordered field of rational numbers, and let  $\mathcal{C}$  be the set of all Cauchy sequences with values in  $\mathbb{Q}$ . More precisely,  $(a_n)_{n=1}^\infty \in \mathcal{C}$  iff  $a_n \in \mathbb{Q}$  for all  $n \in \mathbb{Z}_+$ , and

$$\forall \varepsilon \in \mathbb{Q}, \varepsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N, -\varepsilon < a_m - a_n < \varepsilon.$$

- (a) Define a relation  $R$  on  $\mathcal{C}$  by setting  $(a_n)R(b_n)$  iff  $\lim_{n \rightarrow \infty} (a_n - b_n) = 0$ . Prove that  $R$  is an equivalence relation on  $\mathcal{C}$ .
- (b) Let  $\mathbb{R}$  denote the set of equivalence classes in  $\mathcal{C}$  modulo  $R$ . Define

$$[(a_n)] + [(b_n)] := [(a_n + b_n)], \quad [(a_n)] \cdot [(b_n)] := [(a_n b_n)],$$

for any  $[(a_n)], [(b_n)] \in \mathbb{R}$ . Prove that the above operations of addition and multiplication are well defined (i.e., independent of the choices of representatives of equivalence classes).

- (c) Let  $\varphi : \mathbb{Q} \rightarrow \mathbb{R}$  be defined as  $\varphi(q) = [(q)]$ , where  $(q)$  denotes the constant sequence with all terms equal to  $q$ . Prove that  $\varphi$  is an injection, which preserves the field operations (i.e.,  $\varphi(q_1 + q_2) = \varphi(q_1) + \varphi(q_2)$  and  $\varphi(q_1 q_2) = \varphi(q_1) \cdot \varphi(q_2)$  for all  $q_1, q_2 \in \mathbb{Q}$ .)
- (d) For  $[(a_n)], [(b_n)] \in \mathbb{R}$ , we say that  $[(a_n)] < [(b_n)]$  iff  $\neg([(a_n)] = [(b_n)])$  and there exists  $N \in \mathbb{Z}_+$  such that  $a_n < b_n$  for all  $n \geq N$ . Prove that  $\mathbb{R}$  with so-defined addition, multiplication, and ordering satisfies the axioms of ordered field, in which  $0 = [(0)]$  and  $1 = [(1)]$ . Show that  $q_1 < q_2 \Leftrightarrow \varphi(q_1) < \varphi(q_2)$  for any  $q_1, q_2 \in \mathbb{Q}$  (whence  $\mathbb{R}$  contains  $\mathbb{Q}$  as an ordered subfield).
- (e) Prove that  $\mathbb{Q}$  is everywhere dense in  $\mathbb{R}$ , that is, show that for all  $[(a_n)], [(b_n)] \in \mathbb{R}$ , if  $[(a_n)] < [(b_n)]$  then there exists  $q \in \mathbb{Q}$  such that  $[(a_n)] < [(q)] < [(b_n)]$ , in the above sense.
- (f) **Bonus:** Prove that so-constructed  $\mathbb{R}$  is complete. That is, prove that for every non-empty bounded above set  $X \subset \mathbb{R}$ , there exists  $[(c_n)] \in \mathbb{R}$  such that

$$\forall [(a_n)] \in X, [(a_n)] < [(c_n)]$$

and

$$\forall [(b_n)] \in \mathbb{R} \setminus \{[(c_n)]\}, \quad (\forall [(a_n)] \in X, [(a_n)] < [(b_n)]) \implies [(c_n)] < [(b_n)].$$

**Practice Problems (not to be submitted):**

4. Prove that, if  $P(x)$  and  $Q(x)$  are polynomials of positive degrees, then the sequence  $a_n = \sqrt[n]{\left|\frac{P(n)}{Q(n)}\right|}$  converges to 1. [Hint: You may apply the Algebraic Limit Theorem and Squeeze Theorem, as well as other results proved in class, as needed.]
5. Exercise 5.17.
6. Exercise 5.18.
7. Exercise 6.8.
8. Exercise 6.11.
9. Exercise 6.13.
10. Exercise 6.19.
11. Let  $(a_n)_{n=1}^{\infty}$  be a sequence defined recursively as follows:  $a_1 = 2$ , and  $a_{n+1} = 2 - \frac{1}{a_n}$ , for all  $n \geq 1$ . Prove that the sequence converges and find its limit.
12. Let  $(a_n)_{n=1}^{\infty}$  be a sequence defined recursively as follows:  $a_1 = \sqrt{2}$ ,  $a_{n+1} = \sqrt{2 \cdot a_n}$ , for all  $n \geq 1$ . Prove that the sequence converges and find its limit.
13. Let  $(a_n)_{n=1}^{\infty}$  be a bounded sequence, and suppose that  $\liminf a_n = \limsup a_n$ . Prove that  $(a_n)_{n=1}^{\infty}$  is convergent and  $\lim_{n \rightarrow \infty} a_n = \liminf a_n$ .
14. Let  $(a_n)$  and  $(b_n)$  be two bounded sequences.
  - (a) Prove that  $\liminf a_n + \liminf b_n \leq \liminf(a_n + b_n) \leq \limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$ .
  - (b) Give an example of sequences  $(a_n)$  and  $(b_n)$  for which the leftmost inequality in part (a) is strict.
  - (c) Give an example of sequences  $(a_n)$  and  $(b_n)$  for which the rightmost inequality in part (a) is strict.